


AD-A148 401


INHIBITION OF BABOON MARROW CFU-GEHM CFU-GM BFU-E AND
CFU-E BY ADRENOCOR. (U) BOSTON UNIV MA SCHOOL OF
MEDICINE R ZIEGELSTEIN ET AL. 12 OCT 83 BUSM-83-15
N00014-79-C-0168 F/G 6/15

1/1

UNCLASSIFIED

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

ADA140401

OFFICE OF NAVAL RESEARCH
CONTRACT N00014-79-C-0168

12

TECHNICAL REPORT 83-15

INHIBITION OF BABOON MARROW CFU-GEMM, CFU-GM, BFU-E and CFU-E

BY ADRENOCHROME, AN EPINEPHRINE METABOLITE

by

R. C. ZIEGELSTEIN, S. HOLDEN, A. A. HORLAND, M. D. ALTSCHULE,

AND

C. R. VALERI

NAVAL BLOOD RESEARCH LABORATORY
BOSTON UNIVERSITY SCHOOL OF MEDICINE
615 ALBANY STREET
BOSTON, MA 02118

12 October 1983

Reproduction in whole or in part is permitted for
any purpose of the United States Government

Distribution of this report is unlimited.

DTIC
ELECTE
S D
APR 25 1984
D

DTIC FILE COPY

84 04 24 000

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NBRL, BUSH 83-15	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) INHIBITION OF BABOON MARROW CFU-GEMM, CFU-GM, BFU-E AND CFU-E BY ADRENOCROME, AN EPINEPHRINE METABOLITE		5. TYPE OF REPORT & PERIOD COVERED Technical Report
7. AUTHOR(s) Roy C. Ziegelstein, Sylvia Holden, Allan A. Horland, Mark D. Altschule, and C. Robert Valeri		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Blood Research Laboratory Boston University School of Medicine 615 Albany St., Boston, MA 02118		8. CONTRACT OR GRANT NUMBER(s) N00014-79-C-0168
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Medical Research and Development Command Bethesda, MD 20814		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Naval Medical Command Department of the Navy Washington, D. C. 20372		12. REPORT DATE 12 October 1983
		13. NUMBER OF PAGES 22
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release and sale. Distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Bone Marrow	BFU-E	Hematopoietic stem cells
Adrenochrome	CFU-E	Colony stimulating
Tissue culture	Clonal cell assays	activity
CFU-GEMM	Hemin	Burst-promoting activity
CFU-GM	Red cell production	Blood
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Using tissue culture systems designed for the growth of human bone marrow cells, studies were done to assess the effects of adrenochrome, a metabolite of epinephrine, and of hemin on baboon bone marrow CFU-GEMM, CFU-GM, CFU-E and Bfu-E. In all the assay systems studied, growth of CFU-GEMM, CFU-GM, CFU-E and Bfu-E was suppressed by the addition of 0.5 mM adrenochrome. For those clonal cell assays expressing erythroid growth in culture, the degree of suppression by 0.5 mM adrenochrome increased with increasing cellular maturity		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

(CFU-E > BFU-E > CFU-GEMM).

The addition of 0.2 mM hemin alone produced a slight but not statistically significant depression of baboon marrow BFU-E growth. The combination of hemin and 0.5 mM adrenochrome produced no growth of BFU-E, but did produce a significant increase in the CFU-GEMM as compared to 0.5 mM adrenochrome alone.

These data suggest that substances such as adrenochrome may be involved in the suppression of red cell production observed in patients subjected to traumatic injuries.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ABSTRACT

Using tissue culture systems designed for the growth of human bone marrow cells, studies were done to assess the effects of adrenochrome, a metabolite of epinephrine, and of hemin on baboon bone marrow CFU-GEMM, CFU-GM, CFU-E and BFU-E. In all the assay systems studied, growth of CFU-GEMM, CFU-GM, CFU-E and BFU-E was suppressed by the addition of 0.5 mM adrenochrome. For those clonal cell assays expressing erythroid growth in culture, the degree of suppression by 0.5 mM adrenochrome increased with increasing cellular maturity, (CFU-E > BFU-E > CFU-GEMM).

The addition of 0.2 mM hemin alone produced a slight but not statistically significant depression of baboon marrow BFU-E growth. The combination of hemin and 0.5 mM adrenochrome produced no growth of BFU-E, but did produce a significant increase in the CFU-GEMM as compared to 0.5 mM adrenochrome alone.

These data suggest that substances such as adrenochrome may be involved in the suppression of red cell production observed in patients subjected to traumatic injuries.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
ALL	



INTRODUCTION

Clinicians have long observed that seriously wounded military and civilian persons require larger volumes of blood to restore their blood volumes to normal than can be explained merely by their blood loss.¹⁻⁵ The cause of the unexpectedly low blood volumes, of the so-called "missing blood syndrome", has intrigued many workers who have sought to discover the mechanism for this unexplained hypovolemic anemia. One of the many speculations suggests that circulating toxic substances are involved which inhibit red blood cell production.¹⁻³

In studies by Valeri and Altschule from 1968 to 1972 of more than 300 patients who had sustained war-related injuries in South Vietnam, a clinical pattern of hypovolemic anemia was observed that was characterized by reduced red cell and plasma volumes and normal or slightly reduced peripheral blood hematocrit values and hemoglobin concentration levels.⁶ The reduction in red cell volume was due to blood loss, to a reduction in red cell survival, and to suppression of red cell production. This syndrome was identified as "the hypovolemic anemia of trauma". Because plasma epinephrine concentrations were known to rise above the normal range following traumatic shock,⁷ these authors investigated the possible suppressive effects of epinephrine and its metabolites. Adrenochrome (2,3-dihydro-3-hydroxy-N-methylindole-5,6-quinone), a metabolite formed in the oxidation of epinephrine, was observed in the blood from 25% of the patients with the hypovolemic anemia of trauma and was found to produce hemolysis in vitro.⁸ Toxic substances produced from plasma catecholamines

liberated during traumatic injury were also thought to inhibit erythropoiesis.

We now have available clonal cell assays for pluripotential hematopoietic stem cells producing colonies containing granulocytic, erythroid, monocyte-macrophage, and megakaryocytic elements in methylcellulose (CFU-GEMM),⁹⁻¹¹ for granulocyte-macrophage progenitors in agar (CFU-GM),^{12,13} and for erythroid progenitors in plasma clot (BFU-E, CFU-E),^{14,15} making it possible to study the alterations in blood cell production caused by substances liberated during traumatic injuries.

The purpose of this study was to assess the effect of adrenochrome on baboon marrow CFU-GEMM, CFU-GM, BFU-E and CFU-E. Hemin, the chloride form of hematin (ferric chloride protoporphyrin IX), has been shown to regulate the synthesis of hemoglobin.¹⁶ It has also been shown to enhance the in vitro growth of primitive murine erythroid progenitors.¹⁷ Since red blood cell destruction in the presence of catecholamine metabolites may produce porphyrins and iron,^{6,18} the effects of hemin on the hematopoietic progenitor cells, both alone and in conjunction with adrenochrome, were studied.

MATERIALS AND METHODS

Marrow Samples

Each normal baboon was anesthetized with ketamine hydrochloride, and a 3-5 ml volume of bone marrow aspirated from posterior iliac crests was drawn into heparinized 20 ml syringes which were then stored on ice. Each sample was diluted with an equal amount of Iscove's modified Dulbecco's medium (ImDm) (GIBCO), layered over a ficoll-sodium metrizoate gradient (Lymphoprep, Nyegaard), and centrifuged at 800 X g for 30 minutes at 0-4 C. The mononuclear cells were collected, washed by dilution with ImDm, and then centrifuged at 500 X g for 10 minutes at 4 C. The supernatant solution was discarded and the cells were washed a second time. The cells were counted using a Coulter counter (Coulter Electronics).

Preparation of Feeder Layers for CFU-GEMM and CFU-GM

Human venous peripheral blood was used as the source of colony-stimulating activity (CSA) and burst-promoting activity (BPA), according to the procedure of Pike and Robinson.¹³ Briefly, blood was collected from normal volunteers into heparinized 50 ml syringes and allowed to stand for 1-2 hours at room temperature. The buffy coat was isolated, and the nucleated cells were counted. Feeder underlays were prepared by adding 1×10^6 cells/ml of McCoy's 5A medium (GIBCO) with 20% fetal bovine serum (Sterile Systems) and 0.5% boiled agar (Difco). Each 1 ml aliquot was pipetted into a 35 mm tissue culture plate (Lux) and allowed to gel at room temperature. The plate then received 1200 rads of gamma irradiation (^{137}Cs , Gammacell).

Preparation of PHA-LCM

2×10^6 human peripheral blood leukocytes/ml were stored in Dulbecco's minimum essential medium (MEM) for 7 days in the presence of 1% (v/v) phytohemagglutinin (GIBCO) and 15% fetal bovine serum at 37 C, 5% CO₂ in a humidified atmosphere, as described by Aye et al.¹⁹

Preparation of Hemin and Adrenochrome Solutions

A 2 mM solution of bovine hemin (Type I, Sigma) was prepared according to the method of Stenzel et al.²⁰ 10.4 mg hemin, 0.4 ml 1N KOH (Fisher), and 2.0 ml of filtered 0.2 M Tris-HCl (pH 7.8) were added to distilled water to a volume of 8 ml. The pH was adjusted to 7.5 and the solution was heated in a boiling water bath for 5 minutes. The 2 ml aliquots of this preparation were frozen at -80 C until the time of experiment. A 5 mM solution of adrenochrome (ICN Pharmaceuticals) was prepared on the day of experiment by adding 8.95 mg of adrenochrome to ImDm to a total volume of 10 ml. The unfiltered solution was agitated for 3 minutes prior to use. Hemin was added to yield a final concentration of 0.2 mM. Adrenochrome was added in the final concentrations of 0.005, 0.05, and 0.5 mM.

CFU-GEMM Culture Conditions

CFU-mixed and CFU-GM colonies were grown in methylcellulose, according to the procedure of Ash et al.¹¹ Bone marrow cells in a concentration of 1×10^5 were suspended in 1% methylcellulose in ImDm supplemented with 30% fetal bovine serum, 5% PHA-LCM, 1 U/ml human urinary erythropoietin (EPO, Toyobo), and 5×10^{-5} M 2-mercaptoethanol (Sigma). Four 1 ml aliquots were

layered over feeder underlays and incubated at 37 C in a 5% CO₂ humidified atmosphere for 14 days.

CFU-GM Culture Conditions

CFU-GM colonies were grown in agar-gel, following the procedure of Pike and Robinson.¹³ Bone marrow cells in a concentration of $2 \times 10^5/\text{ml}$ were suspended in 0.3% agar in McCoy's 5A medium supplemented with 20% fetal bovine serum. Four 1 ml aliquots were layered over feeder underlays and incubated for 14 days.

Examination and Scoring of CFU-GEMM and CFU-GM Cultures

Clusters of 50 or more cells, observed unfixed and unstained under a dissecting microscope, were scored as colonies. CFU-GM colonies contained no erythroid cells, whereas mixed colonies contained both non-erythroid and erythroid elements as indicated by their reddish color. Selected mixed colonies which were scored on this basis were examined and found to contain mature granulocytes, erythrocytes, macrophages, and frequently megakaryocytes. The number of CFU-mixed and CFU-GM colonies were reported in the CFU-GEMM assay.

BFU-E Culture Conditions

BFU-E bursts were grown in clotted bovine plasma according to the method of Axelrad et al.¹⁴ One hour before cell plating, an aliquot of bovine plasma (Flow) was placed in a 37 C incubator to allow cold insoluble fibrinogen to dissolve. Bone marrow cells in a concentration of 0.75 to $1.0 \times 10^6/\text{ml}$ were suspended in ImDm supplemented with 30% fetal bovine

serum, 1.5% beef embryo extract (GIBCO), 1% bovine serum albumin (Sigma), 0.2 mg/ml L-asparagine (GIBCO), and 10% warmed plasma. Eight 0.1 ml aliquots of the final mixture were cultured in microtiter wells (Dynatech) placed in 35 mm tissue culture plates and incubated for 12 days.

CFU-E Culture Conditions

CFU-E colonies were grown in clotted bovine plasma according to the method of Stephenson et al.¹⁵ Bone marrow cells in a concentration of $2-2.5 \times 10^5/\text{ml}$ were plated as above for BFU-E, except that a 0.25 U/ml final concentration of human urinary erythropoietin was used and the cells were cultured for 4 days.

Harvesting, Staining, and Scoring of BFU-E and CFU-E Cultures

Each clot was removed from its microtiter well using a metal microspatula and placed on a glass microscope slide. A strip of filter paper (Fisher) was placed over the slide, and clots were fixed with 5% glutaraldehyde (Ladd) and stained with hematoxylin (Fisher) and benzidine (Kodak). Aggregates of 8-64 benzidine-positive cells with characteristic erythroid morphology observed at a magnification of 250X under oil were scored as CFU-E. Isolated groups of 50 or more nucleated cells, at least one-half of which were benzidine-positive observed at a magnification of 100X under oil, were scored as BFU-E.

Statistics

The paired t-test was used to compare control groups to experimental groups

RESULTS

In our laboratory, colonies were grown using the BFU-E and CFU-E assays from baboon bone marrow mononuclear cells under experimental conditions similar to those used in human and murine systems using 2 U/ml EPO for the BFU-E assay and 0.25 U/ml EPO for the CFU-E assay.

The effect of three concentrations of adrenochrome on the growth of marrow CFU-GEMM, CFU-GM, BFU-E and CFU-E was studied. Adrenochrome in a concentration of 0.5 mM inhibited the growth of all hematopoietic progenitor cells studied (Table 1). The degree of suppression varied among the four assays, with the greatest suppression noted in the most mature erythroid progenitor cell (CFU-E: 2.5% control, $p < 0.005$), and the least in the early mixed hematopoietic cell line (CFU-GEMM: 50.6% control, $p < 0.01$). Adrenochrome in lower concentrations did not significantly inhibit growth of CFU-GEMM, CFU-GM, BFU-E or CFU-E ($p > 0.05$).

The effect of 0.2 mM hemin on the growth of marrow CFU-GM, BFU-E, CFU-E and CFU-GEMM and CFU-mixed using the CFU-GEMM assay was also studied. Hemin had no effect on the growth of CFU-GM or CFU-E when compared to untreated controls. In addition, hemin had no effect on these assays in combination with adrenochrome when compared to samples treated with adrenochrome alone (Figures 1 and 2). In contrast, CFU-mixed colonies in the CFU-GEMM assay showed a slight but not statistically significant ($p > 0.05$) increase in number in the presence of hemin alone (Figure 3). In the presence of hemin and 0.5 mM adrenochrome, significant enhancement of mixed colony growth was observed ($p < 0.05$) compared to adrenochrome

FIG.
FIG.

FIG.

alone. At lower adrenochrome concentrations, this effect was not statistically significant ($p > 0.05$).

Growth of BFU-E colonies on day 12 appeared to be suppressed by hemin (Figure 4), but this effect was not statistically significant ($p > 0.2$). However, in the presence of hemin and 0.5 mM adrenochrome, no BFU-E growth was observed in any of the experiments.

FIG. 4

DISCUSSION

Maintenance of mammalian homeostasis depends on the differentiation of several classes of hematopoietic progenitors which appear to arise from the uncommitted stem cell. Among these classes is the pluripotent stem cell which is capable of producing mature granulocyte, erythrocyte, monocyte-macrophage, and megakaryocyte elements in culture (CFU-GEMM), and the CFU-GM, BFU-E, and CFU-E which are derived from it. Our data demonstrate that colonies derived from these four classes of hematopoietic progenitors were grown in vitro from baboon bone marrow mononuclear cells in tissue culture systems designed for the growth of human cells. The number of baboon bone marrow BFU-E colonies on day 12, CFU-E colonies on day 4, CFU-GM colonies on day 14, and CFU-GEMM colonies on day 14, observed in our laboratory were within the range of values reported from other laboratories using human and murine bone marrow.^{17,21-23} This study assessed the effects of adrenochrome, a catecholamine metabolite, and hemin, both alone and in combination, on baboon marrow hematopoietic progenitor cells.

The suppression of colony growth from CFU-GM, CFU-GEMM, BFU-E and CFU-E by the addition of 0.5 mM adrenochrome was observed in all four assay systems studied, although the degree of suppression varied among the assays. In the clonal cell assays capable of expressing erythroid growth in culture, the degree of suppression by 0.5 mM adrenochrome increased with increasing cellular maturity (CFU-E > BFU-E > CFU-GEMM).

Earlier studies on the incubation of adrenochrome and other catecholamine

metabolites in human blood plasma have shown that plasma soluble rheomelanins form after 24 hours at 37 C.²⁴ These rheomelanins have strong free radical activity,^{25,26} which appears to be implicated in red blood cell destruction⁸ and which may be damaging to red blood cell production as well. Free radical reactions are known to disrupt cellular metabolism by inactivating enzymes, cross-linking DNA, and altering membrane functions.²⁷ More recent work has indicated that rheomelanins exhibit several characteristics similar to lipofuscins.²⁸ Therefore, it seems likely that in our study several chemical changes involving adrenochrome occurred in the course of the 4- to 14-day incubation periods.

Although the concentrations of adrenochrome employed were certainly neither pharmacologic nor pathophysiologic, the inhibition of colony growth in vitro suggests a possible role of adrenochrome or one of its metabolites in the anemia of trauma.

Previous observations on the effect of hemin on in vitro growth of murine marrow erythroid progenitors have indicated that hemin in a concentration of 0.2 mM augments the growth of murine marrow BFU-E,¹⁷ but our findings with baboon marrow are in contrast to those observations. Growth of baboon bone marrow BFU-E showed a slight, but not statistically significant, suppression by the addition of hemin. This did not appear to be a non-specific cytotoxic effect on baboon hematopoietic progenitors, since growth of CFU-GEMM was slightly although not significantly enhanced, and growth of CFU-GM and CFU-E was unaffected by the presence of hemin. The reasons for the discrepancies in these results are not clear. They may be due simply to some inherent difference between baboon and murine

marrow hematopoietic cells and their responses to bovine hemin.

Adding hemin to cultures of CFU-GEMM incubated with 0.5 mM adrenochrome increased the colony numbers, an interesting observation insofar as no BFU-E growth was observed in the presence of this same combination. One difference between the two systems is the presence of BPA. Colony growth in the CFU-GEMM assay using methylcellulose contained sources of BPA in the human feeder underlays and in the PHA-LCM, whereas colony growth in the BFU-E assay using plasma clot did not. Since enhanced growth of CFU-GEMM but not BFU-E colony growth occurred in the presence of hemin, it is possible that hemin has no stimulatory effect on baboon marrow erythropoietic progenitors per se, but only brings about an effect through stimulation of human peripheral blood cells to produce increased amounts of BPA. The role of hemin, both alone and in combination with adrenochrome, warrants further study.

FIGURE 1

The effect of adrenochrome on CFU-GM colony formation in agar gel in the presence or absence of 0.2 mM hemin. Values are expressed as the number of CFU-GM colonies per 10^5 marrow cells plated and represent the mean of 5 experiments \pm SEM. No significant difference between cultures with and without hemin ($p > 0.05$).

FIGURE 1

ZIEGELSTEIN ET AL

CFU-GM




FIGURE 2

The effect of adrenochrome on CFU-E colony formation in plasma clot in the presence or absence of 0.2 mM hemin. Values are expressed as the number of CFU-E per 10^5 marrow cells and represent the mean of 5 experiments \pm SEM.




FIGURE 2
ZIEGELSTEIN ET AL

FIGURE 3

The effect of adrenochrome on CFU-GEMM-CFU-MIXED colony formation in methylcellulose in the presence or absence of 0.2 mM hemin. Values are expressed as the number of CFU-GEMM per 10^5 marrow cells and represent the mean of 5 experiments \pm SEM, except where otherwise noted. No significant difference between cultures with and without hemin except in the presence of 0.5 mM adrenochrome ($p < 0.05$). † = mean of 4 experiments.

CFU-GEMM — CFU-Mixed




FIGURE 3

ZIEGELSTEIN ET AL

FIGURE 4

The effect of adrenochrome on BFU-E colony formation in plasma clot in the presence or absence of 0.2 mM hemin. Values are expressed as the number of BFU-E per 10^5 marrow cells and represent the mean of 5 experiments \pm SEM. No significant difference between cultures with and without hemin.

BFU-E




FIGURE 4

TABLE 1

EFFECT OF ADRENOCROME ON CFU-GEMM, CFU-GM, BFU-E and CFU-E*

COLONIES/10⁵ MARROW CELLS

Concentration of Adrenochrome (mM)	CFU-GEMM			BFU-E	CFU-E
	CFU-GM	MIXED	GM		
0	22.9 ± 6.8	15.6 ± 6.4	31.4 ± 8.4	20.6 ± 6.9	353.2 ± 52.4
.005	20.8 ± 6.6**	14.8 ± 7.0**	36.2 ± 12.0**	16.6 ± 7.7**	327.7 ± 52.6**
.05	20.7 ± 6.5**	15.0 ± 5.5**	28.5 ± 6.0**	19.6 ± 6.2**	294.9 ± 45.3**
.5	2.7 ± 1.6 [†] (11.8%)	7.9 ± 5.5 [‡] (50.6%)	5.4 ± 2.9 ^{††} (17.2%)	8.0 ± 6.6 ^{††} (38.8%)	8.8 ± 7.8 ^{†††} (2.5%)

*Data presented as the mean of 5 experiments ± SEM; paired t-tests were done between samples with and without adrenochrome. Percent control values for the 0.5 mM adrenochrome groups appear in parentheses.

**No significant difference ($p > 0.05$)

[†]Significant difference ($p < 0.05$)

[‡]Significant difference ($p < 0.01$)

^{††}Significant difference ($p < 0.025$)

^{†††}Significant difference ($p < 0.005$)

REFERENCES

1. Seamen, B.W. and Ponder, E. The estimation and control of post-operative dehydration with the aid of hemoglobin and plasma protein determinations. *J. Clin. Invest.* 22:673, 1943.
2. Vaughan, J., Thomson, M., and Dyson, M. The blood picture and plasma protein level following injury. *J. Pathol. Bacteriol.* 58:749, 1946.
3. Vaughan, J. Anaemia associated with trauma and sepsis. *Brit. Med. J.* 1:35, 1948.
4. Norlander, O. Total hemoglobin, blood volume and circulatory changes in surgical patients. *Acta Chir. Scand. Suppl.* 199:5, 1955.
5. Topley, E. and Clarke, R. The anemia of trauma. *Blood* 11:357, 1956.
6. Valeri, C.R. and Altschule, M.D. *Hypovolemic Anemia of Trauma: The Missing Blood Syndrome*, CRC Press, Boca Raton, FL, 1981.
7. Benedict, C.R. and Grahame-Smith, D.G. Plasma noradrenaline and adrenaline concentrations and dopamine B-hydroxylase activity in patients with shock due to septicaemia, trauma, and haemorrhage. *Quart. J. Med.* 47:1, 1978.
8. Valeri, C.R., Altschule, M.D., and Pivacek, L.E. The hemolytic action of adrenochrome, an epinephrine metabolite. *J. Med.* 3:20, 1972.
9. Fauser, A.A. and Messner, H.A. Granuloerythropoietic colonies in human bone marrow, peripheral blood and cord blood. *Blood* 52:1243, 1978.

10. Fauser, A.A. and Messner, H.A. Identification of megakaryocytes, macrophages and eosinophils in colonies of human bone marrow containing neutrophilic granulocytes and erythroblasts. *Blood* 53:1023, 1979.
11. Ash, R.C., Detrick, R.A., and Zanjani, E.D. Studies of human pluripotential hemopoietic stem cells (CFU-GEMM) in vitro. *Blood* 58:309, 1981.
12. Bradley, T.R. and Metcalf, D. The growth of mouse bone marrow cells in vitro. *Aust. J. Exp. Biol. Med. Sci.* 44:287, 1966.
13. Pike, B.L. and Robinson, W.A. Human bone marrow colony growth in agar-gel. *J. Cell. Physiol.* 76:77, 1970.
14. Axelrad, A.A., McLeod, D.L., Shreeve, M.M., and Heath, D.S. Properties of cells that produce erythrocytic colonies in vitro. *Hemopoiesis in Culture*, W.A. Robinson, Ed., DHEW Publication No. NIH 74-205:226, 1974.
15. Stephenson, J.R., Axelrad, A.A., McLeod, D.L., and Shreeve, M.M. Induction of colonies of hemoglobin-synthesizing cells by erythropoietin in vitro. *Proc. Nat. Acad. Sci. USA* 68:1542, 1971.
16. Deutsch, J. and Blumenfeld, O.O. Effect of hemin, sodium periodate and concanavalin A on in vitro biosynthesis of rabbit reticulocyte membrane proteins. *Bioc. Biop. Res. Commun.* 79:1224, 1977.
17. Monette, F.C. and Holden, S.A. Hemin enhances the in vitro growth of primitive erythroid progenitor cells. *Blood* 60:527, 1982.

18. Veech, R.L., Rogeness, G.A., and Weil-Malherbe, H. Formation of protoporphyrin from haemoglobin in vitro. *Biochem. J.* 105:1209, 1967.
19. Aye, M.T., Niho, Y., Till, J.E., and McCulloch, E.A. Studies of leukemic cell populations in culture. *Blood* 44:205, 1974.
20. Stenzel, K.H., Rubin, A.L., and Novogrodsky, A. Mitogenic and co-mitogenic properties of hemin. *J. Immunol.* 127:2469, 1981.
21. Ogawa, M., MacEachern, M.D., and Avila, L. Human marrow erythropoiesis in culture: II. Heterogeneity in the morphology, time course of colony formation, and sedimentation velocities of the colony-forming cells. *Am. J. Hematol.* 3:29, 1977.
22. Lipton, J.M., Kudisch, M., and Nathan, D.G. Response of three classes of human erythroid progenitors to the absence of erythropoietin in vitro as a measure of progenitor maturity. *Exp. Hematol.* 9:1035, 1981.
23. Gregory, C.J. and Eaves, A.C. Human marrow cells capable of erythropoietic differentiation in vitro: Definition of three erythroid colony responses. *Blood* 49:855, 1977.
24. Hegedus, Z.L. and Altschule, M.D. Studies on aminochromes: III. Transformation of epinrphrine, adrenochrome and adrenolutin into plasma-soluble melanins during incubation in human blood plasma. *Arch. Biochem. Biophys.* 126:388, 1968.
25. Van Woert, M.H. Reduced nicotinamide-adenine dinucleotide oxidation by melanin: Inhibition by phenothiazines. *Proc. Soc. Exp. Biol. Med.* 129:165, 1968.

26. Mason, H.S., Ingram, D.J.E., and Allen, B. The free radical property of melanins. *Arch. Biochem. Biophys.* 86:225, 1960.
27. DeSesso, J.M. Cell death and free radicals: A mechanism for hydroxy-urea teratogenesis. *Med. Hypoth.* 5:937, 1979.
28. Hegedus, Z.L., Kuttub, S.H., and Altschule, M.D. Studies on rheomelanins: VI. The apparent lipofuscin characteristics of rheomelanins. *Arch. Int. Phys. Bioch.* 88:265, 1980.

END

FILMED

6-84

DTIC